



Microbial N cycling functional gene abundance in forage grazed pasture soils and links to N₂O emissions

Tram Thai, Jiancan Liu, Jesse Reimer, Gazali Issah, Melissa Arcand and Bobbi Helgason
Department of Soil Science, University of Saskatchewan

INTRODUCTION

- Non-bloat legumes (such as Cicer milkvetch (*Astragalus cicer* L.) or Sainfoin (*Onobrychis vicifolia*) are potential alternatives for alfalfa (*Medicago sativa* L.) in forage grazing pastures
- Introduction of the new legumes complicates microbial N cycling and the link to N₂O under these systems.
- Information on N₂O emissions from soil and N functional gene abundance in semi-arid grassland is limited.

OBJECTIVES

- Characterize N functional gene abundance in soils under different grass-legume grazing pasture.
- Investigate the relationships among bloat and non-bloat legumes, N₂O emissions, N-related soil chemical and microbiological parameters.




Diagram 1: Microbial N cycling and key N functional genes in soils under grazed forage legume pasture


MATERIALS AND METHODS

- Termuende Research Farm, Lanigan, SK.
- Established grass/Algonquin Alfalfa stand (Control), sod seeded non-bloat legumes (Common Sainfoin, Veldt Cicer milkvetch) x 3 replicate grazed paddocks (65 x 300m x 4-5 steers/year)
- Soil N₂O flux chambers.
- Soil samples (0-10 cm): 3 regions x 9 paddocks x single timepoint (August) x 2 years (2017, 2018)
- Soil DNA extraction and qPCR analysis.

REGION	Algonquin Alfalfa (A)	Algonquin Alfalfa (A)	Algonquin Alfalfa (A)
Far (F)			
Middle (M)			
Close (C)			
	W	W	W

REGION	Common Sainfoin (S)	Veldt Cicer milkvetch (C)	Common Sainfoin (S)	Veldt Cicer milkvetch (C)	Common Sainfoin (S)	Veldt Cicer milkvetch (C)
Close (C)						
Middle (M)						
Far (F)						
	W	W	W	W	W	W

W: Water source



RESULTS




Figure 2: N functional gene abundance in soils under different grazed legume-grass mixture in August, 2017 and 2018




Figure 3: Correlation among N functional gene abundance, N₂O emissions and N-related soil parameters

Figure 4: RDA biplot among explanatory variables (species, region), N functional gene abundance, N₂O emissions and N-related soil parameters

MAIN FINDINGS, DISCUSSION AND CONCLUSION

- Despite no difference in N₂O emission under different grass-legume mixtures (Fig.1), AOB gene abundance was higher in non-bloat grass pasture soils (Fig.2), indicating a potential nitrification (by bacteria) under these pasture.
- In Cicer milkvetch: There were (i): a *species x region* interaction effect on *NirS* gene abundance, which was more abundant in the far region - the most crowded area; (ii) a *year x species* interaction effect on *NosZl* abundance in 2018-a drier year (Fig.2), which suggest a key role of grazing density and water on these denitrifiers.
- Gene abundance (AOB, *NirS/K*, *NosZl*) was not correlated to N₂O, but AOB and *NirS* were correlated with soil pH (Fig.3).
- Soil pH as well as AOB, *NirS* gene abundance were positively associated with Cicer milkvetch, might be potential predictors for N cycling in soil under this pasture.

ACKNOWLEDGEMENTS

The research was funded by Agricultural and Agri-Food Canada from Agricultural Greenhouse Gases Program (AGGP) project. The authors would gratefully acknowledge the technical support from AGGP project team members (lab 5C15, 5E19, 5E25, 5E31), collaborators, as well as support from summer students.

CONTACT INFORMATION

Tram.thai@usask.ca

